General Index

A

Acanthodesia tenuis, 83, 84, 228
Acanthohaustorius intermedius, 187, 188, 189, 191
  millsi, 187, 188, 189, 191, 192
Acartia tonsa, 83, 259, 260
Achromobacter, 146, 153
Acorn barnacle, 77
Aeromonas sp., 143, 147, 148, 153
  formicans, 148, 149
  liquefaciens, 137, 147, 148, 149
  salmonicida, 137, 143, 149
Agar, composition, bacterial cultures, 137
Alewife, 127
Algae, 22, 76
Allen, N. and M. J. Pelczar, Jr., 135–54
American eel, 255
American oyster, 228, 261
Amidostomum, 68
Amphidinium sp., 92, 97
Amphipod, 20, 76, 80, 85
  community structure, 187
  concentrations, 190–1
  density, control factor, 81
  distribution, 187–92
  ecological "niche," 188–9
  egg number, 189–91
  size, female, 189–91
  spawning season, 190–2
  tube building, 80–1
Amphiporeia virginiana, 187, 189, 190, 191
Amphiprora alata, 92, 100
Amphiuma, 22
Amygdalum papyria, 20
Anchoa mitchilli, 128
Anchovy, 127, 128
  spawning patterns, 128
Andrews, J. D. and J. L. Wood, 1–13
Anemones, 76–7, 82–3, 212
Anguilla, 20
Anguilla rostrata, 255
Anodonta imbecilis, 23
Anodontid mussels, 21
Anthurid isopod, 14, 21
Antibody
  Gel diffusion test, 66
  striped bass, 66
  white perch, 66
  sensitivity, 139
Antigen, 66
Aquaria
  clam, experimental, 194
  fish, closed, 143
Arlina, 29
  effulgens, 35, 46
  effulgens = *Etheostoma olmstedi olmstedi*, 29
  maculaticeps, 48
Armandia agilis, 191
Arrowleaf, 15
Artemia, 171, 173, 174
Anthropod
  predator-prey relations, 85–6
Asbestos, surface, fouling organisms, 229–35
Ashby, W. and L. P. Schultz, 237–52
Ashing, organic matter, 254, 262
Astacus sp., 170, 171, 173, 174, 175
Asterias forbesi, 173
Asterionella formosa, 92, 100
  japonica, 90, 92, 93, 94–5, 97, 100
  distribution, 94–5
Atlantic croaker, 129
  silverside, 255
Aurelia aurita, 217

B

Bacillus, 145, 152
Bacteria, 76
Bacterial cultures, 137
  similarity coefficient, 141, 143–5, 147–8, 153
  white perch, 136–8
Bacteriastrum delicatulum, 92, 100
Bacteriological studies, computer, 141–3
Bagley, G. E. and L. N. Locke, 68–9
Bailey, R. L. and C. E. Richards, 204–5
Balanus eburneus, 77, 78
  improvisus, 76, 77, 78, 228
Barnacle, 76–8, 80, 86, 215, 228–9, 231, 233
  acorn, 77
  competition, community, 78
  ivory, 77
  light, exposure to, 78
  predation by flatworm, 78
Bass
  largemouth, 22, 127
  warmouth, 22
Bathyporeia, 192
Bay scallop, 155, 166
Beach
  conservation
    factors, 242
    groin effects, 238–52
  erosion
    factors, 242, 243
    groin construction, 238–9, 242, 247, 250, 251
Beam trawl, 130
Beckett, R. L., 261–2
Bent mussel, 83
Biddulphia longicruris, 92, 100
  mobiliensis, 92, 100
  rhombus, 92, 100
Bimeria franciscana, 76, 79, 80, 81, 84, 86
Biochemical analyses, white perch, 140–1
Biocides, 253
Biological stains, 171–2
Biology, sea nettle, 209–20
Biotic succession, epifauna, 76–7
Bird, blood parasite of, 200
Bishop, J. W., 259–60
Blenny, striped, 127
Blood
  cell morphology, lobster, 170–86
  parasite, bird, 200–1
  sampling, lobster, 171
  types, lobster, 172–3
Blueback herring, 127
  Blue crab, 171, 202
  Bluegills, 263
  Blue shark, 53
  Boleosoma (*Etheostoma*), 29, 30, 31, 34, 35, 41, 43, 47
    canal system, 34
    cephalic system, 34
    description, 29, 30, 34
    distribution, 43
    synonymy, 29
  Boleosoma aesopus, 35
    atromaculata, 41
  camurum, 29
  effulgens, 36, 48
  longimanus, 36
  maculaticeps, 48
  olmstedi, 35, 41, 48
    aesopus, 36
    atromaculatum, 41
    effulgens, 36
    nigromaculata, 39
    olmstedi, 35, 36
    var. *olmstedi*, 36
  nigrum, 36, 41, 46, 48
    effulgens, 36
    nigrum, 36
    olmstedi, 36
    vexillare, 46
  stigmaeum, 29
    tessellatum, 35, 36
    = *Etheostoma olmstedi olmstedi*, 29
  vexillare, 46
Bolinopsis, 259
Bowerbankia gracilis, 83
Brachiodontes recurvus, 83, 228
Brachiura sowerbyi, 22
Branchipus, 171, 173, 174
Branta canadensis, 68
Bream, 21
Broad Creek, Maryland
  description, 228–9
Bryazoan, 77, 78, 83, 215, 228–31, 233
  zooids, 84
Bull shark, 52
Burbanck, G. P. and W. D. Burbanck, 14–27
Burbanck, W. D. and G. P. Burbanck, 14–27

C

Callinectes ornatus, 202
  sapidus, 20, 22, 23, 171, 174, 202, 203
  similis, 202
  spp., 202, 203

Cambarus fallax, 22
Campanularidae, 79, 80
Campostoma anomalum pullum
distribution, 39
Canada geese, 68–9
Carbohydrate fermentation tests, 140
Carcharhinus, 59
leucas, 52
milberti, 52
Carcharodon carcharias, 52
Cargo, D. G. and L. P. Schultz, 209–20
Carunculina minor, 23
Catfish, 22, 86
Cerataulina bergonii, 92, 100
Ceratium furca, 92, 97
fusus, 97
tripos, 92, 97
Chaetoceros affinis, 92, 100
brevis, 92, 100
compressus, 92, 100
decipiens, 92, 100
peruvianus, 92, 100
subtilis, 92
Chara, 22
Chesapeake Bay
fish nursery, 127–8
MSX infections, 3, 5
Chilomonas sp., 97, 100
Chincoteague Bay
fungus occurrence, oysters, 261
Chiridotea caeca, 191
Chironomid, dipterous, 21
Chironomids, 22
Chiropsalmus quadrigatus, 210
Chroomanas vectensis, 92
Chrysaora, 210, 212, 213, 215, 217, 218, 219
medusae, 215
quinquecirrha, 52, 209, 216
Clam, 254–6, 263
activity, temperature, 194–6
aquaria, experimental, 194
density, natural, 196
hydraulic dredge
efficiency, 193
operation, 193
marketable size, 196
metal retention, 263
reburrowing, activity, 193–4, 198
hydraulics, 198
sediment composition, 196–7
size correlation, 198
temperature effects, 194–6
time, 198
Clingfish, 77, 86, 127
predator-prey relations, 85–6
Clogging, plankton net, 213–4
Closterium gracile, 92, 100
Clymenella torquata, 191
Coccidiosis, Canada geese, 68–9
Cocconeis scutellum, 92
Coelenterate
food items, 217
Cole, C. F., 28–51
Columba livia, 200
Community
competition, barnacle, 76–8, 80, 86
structure, amphipods, 187
Computer, bacteriological studies, 141–3
Copepod, 83, 259, 260
density, Patuxent River, 260
mortality rate, estuary, 260
Corambella, 84
Cordgrass, 221
Cordylophora, 224
Cordylophora lacustris, 79, 80, 81, 84
Corethron hystrix, 92, 100
Cornus florida, 264
Corophoriids, 21
Corophium lacustre, 76
Cory, R. L., 71–89
Coryphella, 214, 217
Coscinodiscus asteromphalus, 92
perforatus, 92
Crab, blue, 171, 202
mud, 77, 80, 86
ovigerous females, 203
Crangon septemspinosa, 76, 86, 256
Crassostrea virginica, 1, 79, 85, 228, 233
Crayfish, 22, 170
Croaker, Atlantic, 129
Crows, 22
Cryptomanas sp., 92, 93, 94, 100
distribution, 97
salina, 92
stigmatica, 92
Ctenophore, 217–8, 259–60
feeding rate, 259–60
food, jellyfish, 217, 218
length, 259, 260
oyster predator, 217
predator, 259, 260
respiration, 259
volume, 259
Culicoides, 201
Culture, oyster, 2, 136–8, 141, 143–5, 147–8, 153
Current
drag, estuary, 156–8
estuary, 160
velocities, estuary, 159
Cyanea, 219
capillata, 219
Cyathura, 14–5, 20, 21, 22
biotypes, 15, 16, 20–4
depth requirements, 17, 25
environmental stress, 14–5
habitat, 16, 20–4
oxygen requirements, 14, 16, 21, 22, 24–5
pH requirements, 14, 16, 21, 26
salinity requirements, 14, 16, 25, 26
salinity tolerances, 25
stress, environmental, 14–5
temperature tolerance, 25
carinata, 15, 25–6
seasonal changes, 25–6
polita, 14–7, 20–7
temperature requirements, 14, 16, 20–1, 25–6
Cyclotella striata, 92, 100
Cymadusa compta, 85
Cynoscion regalis, 129
Cyprinid, 39
Cyprinodon variegatus, 67
Cyst, sea nettle, 212, 216–7
Cytochrome oxidase test, 139, 146

D

Daiber, F. C. and R. J. Reimold, 132–3
Damsel fly, 21
Darter, 31
Johnny, 28–50
lateral line scales, 37
Dermocystidium marinum, 3, 7, 11, 261
Dexter, Deborah M., 187–92
Diadumene leucolena, 212
Diatoma hiemale, 92
Diatoms, 90–100
seasonal variation, 90, 91–7, 100
Dinoflagellate, 67–8
Dinophysis sp., 92
Diopatra cuprea, 191
Dipterous chironomid, 21
Dissolved O_2 analyses, estuary, 158
Ditylum brightwellii, 92, 100
Diurnal-nocturnal cycle, interstitial water, 26
Donax variabilis, 191
Doration, 29–30
description, 29–30
Dove, white-winged, 201
Dovel, W., 125–9
Dragonfly, 21, 23
Drobeck, K. G. and H. T. Pfitzenmeyer, 193–9
Ducks, 201

E

E. meridianales, 78
Eastern mourning dove, 200–1
Ecological niche, amphipods, 188–9
Eisler, K., 262–4
Eisler, K. and M. P. Weinstein, 253–8
Eel, 20
Egrets, 22
Elliptio buckleyi, 22, 23
Emerita talpoida, 191
Enterobacter, 149, 153
aerogenes, 149
Environmental stress, *Cyathura*, 14–5
Ephemera simulans, 14, 25
Epibenthic environment, characteristics, estuary, 155–69
Epifauna
biotic succession, 76–7
central factors, 74–5
Patuxent River, 71–89
productivity, estuary, 86–8
salinity, 74
sampling, 72
temperature, 74–5
temporal sequence, 76–7
turbidity, 75
Epizootic, fish, 135
oyster, 1–13
Erosion, beach, 237–52
Estrella, 29
atromaculata, 29, 39
Estuary, copepod density, 260
copepod
mortality, 260

current drag, 156–8
shear, 160
velocities, 159
Estuary
epibenthic environment, 155–69
epifauna, 71–89
eutrophication, 132–3
fluorometric observations, 157
oxygen, dissolved, 158
phosphorous, 132–3
analyses, 157
concentrations, 132–3
sources, 132
radiant energy, 158
sampler, microstratification, 157
silicate analyses, 157
turbidity, 158
Eteone heteropoda, 85
Etheostoma, 29, 34, 49
aesopus, 36
chlorosomum, 29
effulgens, 36, 37, 39
fusiforme barratti, 44
jessiae, 29
longimanum, 29, 47
nigrum, 29, 31, 33, 34, 35, 36, 37, 46, 47, 48, 49
comparison, *E. olmstedi*, 33
description, 33–4
distribution, 43–45
effulgens, 36, 37
description, 37
distribution, 44
lateral line scales, 37
eulepis, 29, 43
distribution, 43
maculaticeps, 49
nigrum, 29, 31, 43, 44–5, 48
olmstedi, 29, 36
vexillare, 36
olmstedi, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41–6, 47, 49
description, 31–5
distribution, 31, 32, 37, 39
glaciers, effects of, 37–39
lateral line scales, 31, 37, 38
salinity requirements, 43–4
subspecies characteristics, 33
olmstedi atromaculatum, 29, 32, 33, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47
characteristics, 33, 36, 41
lateral line scales, 38, 44
salinity requirements, 43–4
spawning, 42
squamation factors, 40, 41–2, 44
synonymy, 39, 41
olmstedi effulgens
description, 37
glaciers, effects of, 37, 44–6
lateral line scales, 37
olmstedi maculaticeps, 29, 31, 32, 33, 36, 37, 38, 40, 48
characteristics, 33, 36, 48–9
distribution, 48–9
lateral line scales, 38, 48
squamation factors, 40
synonymy, 48
olmstedi olmstedi, 29, 32, 33, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47
characteristics, 33, 36, 37
distribution, 32, 39, 44
glaciers, effects of, 37, 44–6
lateral line scales, 37, 38, 39, 44
squamation factors, 39, 40, 41–2, 44
synonymy, 35–36
olmstedi vexillare, 29, 30, 32, 33, 37, 38, 40, 41, 46, 47
characteristics, 33, 36, 46
distribution, 46–7
glaciers, effects of, 47
lateral line scales, 38, 42
squamation factors, 40, 46
synonymy, 46
perlongum, 29, 41
podostemone, 29, 47
sellare, 41
stigmaeum, 29
Eucampia zoodiacus, 92, 100
Euglena sp., 92, 100
Euryhaline gammarid, 22
Euryokous isopod, 26
Eutrophication, estuarine, 132, 133
Exosphaeroma diminutum, 191
Exuviaella sp. 92, 97
marina, 92

F

Facelina droumondi, 217
Fecundity, amphipods, 189–91
Feeding rate, ctenophore, 259–60
Fish
antibiotic sensitivity, 139
antibody, 66
aquaria, closed, 143
bacteria cultures, 136–8
biochemical analysis, 140–1
commercial value, 103
epizootic, 135
homing tendency, 122
lateral line system, 30, 31, 37, 38, 39, 42, 44
mark and recapture, 105–123
metal toxicity, 262–3
meristics, 204
migration, 102–24
nursery, Chesapeake Bay, 127–8, 129
organ-body weight relationship, 63–5
organ growth, 64
pigment production, test for, 138
predator-prey relationship, 85–6
salinity requirements, 43–4
seasonal movements, 102–24
spawning, 42, 127–8, 129
sport value, 103
staining, 138
surface trawl, 130–1
tagging, 104–5
taxonomy, 30
toxicity, zinc, 262–3
vertical distribution, 109
zinc, toxicity, 262–3
Fish kill, Chesapeake Bay, 135
Flagellate, 90, 101
seasonal variations, estuary, 90, 91–7, 100
Flatworm, 77, 78, 228, 229, 231
barnacle predator, 78
Flies, hippoboscid, 201
Flounder, 22
winter, 127, 128
Fluorometric observations, estuary, 157
Fouling organisms
setting, 229–35
surface preferences, 231, 232, 233, 235
Freshwater fish, taxonomy, 30
Fundulus heteroclitus, 67, 255, 262
luciae, 204, 205
majalis, 67
Fungus, 3
oyster, 261

G

Gammarid, 21, 23
Gammarus tigrinus, 22, 23
Gapers, oysters, 2
Geese, 201
Geese, Canada, 68–9
Gellius, 83
Gizzard worms, 68
Glaciers, affecting fish distribution, 37–9, 44–6
Glycera dibranchiata, 191
Gobiosoma bosci, 86
Gobisox strumosus, 86
Goby, 127
Goby, green, 127
Goniobiasis floridanus, 22
Gonyaulax sp., 92, 94, 97
Grammatophora marina, 92
Grass
shrimp, 86, 256
tape, 15, 20
turtle, 15
Gray sharks, 53
Green goby, 127
Greenland shark, 53
Groin
beach obstruction, 238–52
construction, 238–9, 242, 247, 250–1
Guinardia flaccida, 92, 100
Gymnodinium sp., 92, 97
Gyrosigma balticum, 92, 100
spencerii, 92, 100

H

Haemoproteus, 200, 201
columbae, 200, 201
maccallumi, 200, 201
sacharovi, 200, 201
Hammerhead sharks, 52, 53
Haploscoloplos fragilis, 191
Haustoriids, 20, 23
Haustorius arenarius, 192
Haustorius sp., 26, 187, 189, 190, 191
Hearing, V. and S. H. Vernick, 170–86
Hemosiderosis
Canada geese, 69
mallards, 69
Herman, C. M. and J. O. Knisley, Jr., 200–2
Hermit crab, 256

Herons, 22
Herring, 127
  blueback, 127
Hickory shad, 127
Hogchoker, 127
  spawning patterns, 128
Hololepis, 30
Homarus, 171, 173, 174, 175
  americanus, 170, 171, 172
Homing tendency, fish, 122
Hooked mussel, 228
Haplosporidian parasite, 1
Hydraulic dredge, clam, 193
Hydrogen-ion concentration, interstitial water, 26
Hydroids, 76, 77, 78, 79–81, 84, 88
  polyps, 80, 84
Hippoboscid flies, 201
  Haemoproteus, 201

I

Ibis, 22
Idotea baltica, 85
Insecticides, 253–8
Interstitial water
  diurnal cycle, 26
  hydrogen-ion concentration, 26
  oxygen determination, 19, 20, 24–5, 26
  pH
    determinations, 17, 19–20, 26
    diurnal nocturnal cycle, 17, 19–20, 26
  salinity, 17, 19, 26
  sampler, description of, 17, 24
    operation, 17–19, 25
  sampling, 14–27
  temperature, 19, 25–6
Ioa, 30
  description, 30
Isopod, 85
  anthurid, 14, 21
  sampling, 14–27
Ivory barnacle, 77

J

James River, 90, 91, 99–100
  oysters, seed area, 2
  phytoplankton, 90–101
  plankton, 90–101
  salinity range, 90, 99
  surface temperature, 93, 94, 95, 99
Jansen, W. A. and C. D. Meyers, 66–7
Jellyfish
  food, 217–8
  winter, 219
Jellyfishes, 209, 210
Johnny darter, 28–50

K

Killifish, spotfin, 204
Knisley, J. O., Jr. and C. M. Herman, 200–2

L

Lamna carnubica, 57
Largemouth bass, 22, 127
Lateral line
  environmental control, 37
  scales, fishes, 37
  system, fish, 30
Lawler, A. R., 67–8
Lead poisoning, Canada geese, 68–9
Leeches, 23
Leiostomus xanthurus, 129
Lent, C. M., 221–7
Lepidactylus dytiscus, 187, 189, 190, 191
Leptochelia, 85
Leptocylindrus danicus, 92, 93, 94
Leptosynapta inhaerens, 191
Licmophora abbreviata, 92, 100
Limulus, 174, 175
Liquidambar straciflua, 264
Liriodendron tuplifera, 264
Live trap, shrew, 264
Lobster, 170, 172–4
  blood cell
    morphology, 170–86
    types, 172–3
  blood sampling, 171
Locke, L. N. and G. E. Bagley 68–9
Lonicera japonica, 264
Lucania parva, 67

M

Macoma balthica, 85
  mitchelli, 20, 26
Magelona papillicornis, 191
Magothy River
  description, 125–6, 127, 129
  nursery, fish, 129
  salinity, 126
  spawning area, fish, 129
Mako shark, 53, 61
Malathion, insecticide, 253–8
Mallard, 69
Mark and recapture, fish, 105–23
Marshall, H. G., 90–101
Marshall, N., 155–69
Marsh mussels, 224
Mayfly, burrowing, 14, 23, 25
Melita nitida, 85
Mellita quinquiesperforata, 191
Melosira jurgensii, 92, 100
  sulcata, 92, 93, 100
Membranipora crustulenta, 83, 228
  membranacea, 83, 84
Menhaden, 127, 128
Menidia menidia, 255
Mercenaria mercenaria, 191, 197, 253
Metal
  clam
    accumulation, 254–7
    retention, 263
Methoxychlor, insecticide, 253–8
Meyers, C. D. and W. A. Janssen, 66–7
Microcionia prolifera, 83
Micrococcus, 150, 153
Micropogon undulatus, 129
Migration
  fish, 105–24
  striped bass, 102, 124
Miller, R. V. and P. R. Nichols, 102–24
Minchinia nelsoni, 1
Mnemiopsis leidyi, 217, 259, 260
Modiolus, 225, 226
  (*Arcuatula*) *demissus*, 221, 222, 223, 225
Molgula manhattensis, 77, 78–9
Molluscs, 225, 226
Morphometrics, ribbed mussel, 222–6
Mortality, copepod, estuary, 260
Mourning dove, eastern, 200, 201
MSX (multiple spheroid X), 1–3, 5–13
  activity variations, 12
  Chesapeake Bay occurrence, 3, 5
  distribution, 1, 2, 3, 5, 6–7, 10
    Virginia, 5, 12
  epizootic, 3
  history, Virginia, 3–5
  infections, initial, Chesapeake Bay, 3, 5
  intensities, annual variations, 2, 10, 12
  mortality patterns, Virginia, 4–5
  mortalities, temperature induced, 3
  origin, Chesapeake Bay, 11
  oyster reduction, 3
  salinity range, 1
  type areas, 2–3, 7–9, 10
Mud crabs, 77, 80, 86
Mugil cephalus, 255
Mummichog, 255, 262, 263
Musculium, 21
  transversum, 23
Mussel, 3, 21, 22, 23, 229, 230, 233
  bent, 83
Mya arenaria, 196, 198, 221, 224
  edulis, 221
Mytilid, 20
Mytilus, 225, 226
  californianus, 221, 224
  edulis, 3
    setting time, 3

N

Nassarius obsoletus, 191
  vibex, 191
Navicula sp., 92
Needlefish, 127
Neohaustorius schmitzi, 187, 188, 191
Nephtys picta, 191
Nereids, 20, 21, 23
Nereis succinea, 85
Neretina, 20
Neriid worms, 77
Niche, ecological, amphipods, 188–9
Nichols, P. R. and R. V. Miller, 102–124
Nitrate analyses, estuarine water, 157
Nitzschia closterium, 92, 100
  longissima, 92, 100
  paradoxa, 92, 100
  pungens atlantica, 90, 92, 93, 94, 96, 100
    distribution, 96
  seriata, 92, 93, 94, 96, 97, 100
    distribution, 96
Nocturnal-diurnal cycle, interstitial water, 26

Northern puffer, 255
Nudibranch, 81, 84–5, 214, 217
as sea nettle predator, 214

O

Ocypode albicans, 191
Oligocephalus, 30
description, 30
Oligochaete worms, 21, 22, 23
Oodinium cyprinadontum, 67, 68
cyst formation, 68
description, 67–8
dinospore formation, 68
parasitic stage, 67–8
range, 67
poucheti, 68
description, 68
Ophiuroids, 22
Orchestia platensis, 191
Organic matter, asking, 254, 262
Oscillatoria sp., 92, 100
Otter trawl, 130
Oxygen
cyathura
requirements, 14, 16, 21, 22, 24–5
interstitial water, 19, 20, 24–5
determination, 19, 20, 24–5
Oxyurostylis smithi, 191
Oyster, 1–13, 78, 83, 85, 229–33, 235
culture, 2
epizootic, 1–13
fungus, 261
gapers, 2
growing areas, Virginia, 2–3
James River seed area, 2
mortality, 1–13
Virginia (1958–9), 3, 5
pathogen, 1–13
plantings, 11–12
population reduction, MSX, 3
seed area, 2
spat, 233
worm, 85

P

Pagurids, 22
Pagurus longicarpus, 256
Palaemonetes vulgaris, 76, 86, 256
Parahaustorius longimerus, 187, 188, 189, 191
Pathogen, oysters, 1–13
Pathogenicity, white perch bacteria, 136, 142, 150
Patuxent River
epifauna, 71–89
productivity, 86–8
Pelczar, J. J., Jr. and M. Allen, 135–54
Perca flavescens, 128
minima, 35
Perch
white, 66, 127, 135
yellow, 126, 128
Percid, 44
Percina peltata, 39
Peridinium sp., 92
Peridinium triquetrum, 97
Pesticide, induced stress, 253–8
Pfitzenmeyer, H. T. and K. G. Drobeck, 193–9
pH
Cyathura, 14, 16, 21, 26
requirements, 14, 16, 21, 26
determinations, 17, 19–20
interstitial water, 26
Phosphorous
estuarine, 132–3
analyses, 157
Phragmites, 22
Phytoplankton
James River, 90–101
vertical distribution, 97
Pigeon, 200, 201
Pigment production, test for, fish, 138
Pinus taeda, 264
Pipefish, 127
"Pit" trap, shrew, 264
Plankton, estuary
distribution
seasonal, 91–7
vertical, 97–9
James River, 90, 101
nets, clogging, 43–4, 213–4
sampling, 259
Planktoniella sol, 92, 100
Planulae, sea nettle, 214–5
Pleurosigma angulatum, 92, 100
formosum, 92, 100
Polychaetes, 22
Polydora ligni, 81–2, 85
Polyp, sea nettle, 210, 212, 214, 215
Porifera, 83
Predator-prey relationship
fish-amphipod, 85–6
flatworm-barnacle, 78
nudibranch-sea nettle, 214
sea nettle-ctenophore, 217–8
Pristina osborni, 22
Productivity
epifauna, 86–8
estuary, 86–8
Patuxent River, 86–8
Prorocentrum sp., 94, 97
micans, 92, 97
minimum, 92
triangulatum, 92
Protozoans, 76, 83
Protohaustorius deichmannae, 187, 188, 191
Pseudomonas sp., 137, 143, 146, 153
aeruginosa, 145
Pseudopleuronectes americanus, 128
Puffer, fish, 63–5
distribution, 63
organ-body weight ratios, 63–5
organ growth, 64
brain, 64
heart, 64
heart, size determination, 65

Q

Quahaugs, 253, 255, 256, 257
mantle, function, 258
metal accumulation, 254–7
Quercus sp., 264

R

Radiant energy determination, estuary, 158
Rangia cuneata, 20, 21
Ranunculus, 224
Reburrowing, clam, 193–4, 198
Recaptures marking, fish, 105–23
Red sponge, 83
Reimold, R. J. and F. C. Daiber, 132–3
Respiration, ctenophore, 259
Rhasphoneis amphiceros, 92
Rhithropanopeus, 22, 23
Rhizosolenia alata, 92, 93, 96–7, 100
distribution, 96–7
calcar avis, 92, 100
delicatula, 92, 100
fragilissima, 92, 93, 94, 100
setigera, 92, 93, 97, 100
distribution, 97
stolterfothii, 92, 93, 96, 100
distribution, 96
Rhopilema verrilli, 217
Ribbed mussel, 221–3
growth indices, 221–7
habitat effects, 221–7
morphometrics, 222–6
shell morphology, 222–3
shell-weight/meat-weight, 221
Richards, C. E. and R. L. Bailey, 204–5
Rithropanopeus harrisi, 20, 22, 86
Roccus americanus, 66, 127, 135
saxatilis, 66, 102–24, 127
Ruppia maritima, 83

S

Sagartia leucolena, 76, 82–3
Sagittaria, 15
Salamander, 22
Salinity
Cyathura
requirements, 14, 16, 25–6
darter
requirements, 43–4
determinations, 17, 19
interstitial waters, 26
epifauna, 74
interstitial water, 17, 19, 26
James River, 90, 99
Magothy River, 125, 126, 127, 129
MSX range, 1
range, polyp, sea nettle, 210
tolerance, jelly fish, 219
sea nettle, 218
Salmonids, 263
Sampler
interstitial water
description, 17, 24
operation, 17–9, 25
microstratification, estuary, 157
Sampling
epifauna, estuarine, 72
interstitial water, 14–27
isopods, 14–27
plankton, estuary, 259
surface trawl, fish, shrimp, 130–1
Sand shrimp, 86, 256
Scallop, bag, 155, 166
Schneider, R. and C. G. Wilber, 63–5

Schultz, L. P., 52–62
and W. Ashby, 237–52
and D. G. Cargo, 209–20
Sciaenids, 129
Scientists Cliffs, Md.
sediment composition, 237
Scolelepis squamata, 191
Seals, 53
Sea nettle, 52, 209, 212
biology, 209–20
diameter, 212
medusa size, 213
pedal cyst, 212, 216, 217
planulae, setting, 214, 215
polyp distribution, 210
polyp growth, 212, 214, 215
polyp salinity range, 210
salinity tolerance, 218
strobilation
period, 211, 214
temperature, 214, 215
temperature tolerance, 216–7
treatment, 210–1
Sea slugs, 84, 217
Seasonal movements, fish, 102–24
Seasonal variation, diatoms, 90, 91–7, 100
Sea squirt, mortalities, 79
Sea wasp, 210
Sediment
Chesapeake Bay, 237
composition, clam habitat, 196–7
fishes, effects on, 37–9
glacial, past, 37–9, 44–6
Seine, 202
fish, 204
Setting
fouling organisms, 228–36
asbestos plates, 229–35
Shad, hickory, 127
Shark
attacks, 52–62
conditions for, 52, 53, 56, 59, 61
prevention, 56, 57–8, 60, 61
blue, 52, 53
gray, 53
Greenland, 53
hammerhead, 52, 53
mako, 53, 61
tiger, 53, 61
white, 52, 53, 60, 61
white-tip, 53
vision, 57
Shaw, W. N., 228–36
Shear influences, current, estuary, 160
Shrew, 264
live trap, 264
Virginia occurrence 264
shrimp, 76
grass, 86
sand, 86
trawl, 130–1
Silicate analyses, estuary, 157
Silversides, 127, 129
vertical distribution, 129
Similarity co-efficient, bacterial cultures, 141, 143, 144, 145, 147, 148, 153
Skates, 22
Skeletonema costatum, 90, 92, 93, 94, 95–6, 97, 100, 101
distribution, 95–6, 100–01
Slug, sea, 84, 217
Snail, 21, 22
Snapper, 61
Soft-shelled clams, 221
Solemya velum, 191
Somniosus microcephalus, 53
pacificus, 53
Sorex longirostris, 264
Spartina, 20, 221
Spawning
amphipod, 190–2
anchovy, 128
fish, 42, 128
Sphaeroides maculatus, 63, 64, 255
Sphaeromids, 23
Sphyrna tiburo, 52
Sponges, 215
red, 83
Spot, 129
Spotfin killifish, 204
meristics, 204
ova, 204–5
Staining
blood cells, lobster, 171
fish, 138
Stains, biological, 171–2
Starfish, 173
Stephanopyxis turris, 92
Stizostedion marinum, 44
Stout, J., 264
Stress
Cyathura, environmental, 14–5
pesticide-induced, animal, 253–8
Striped bass, 23, 102–24, 127
antibody, 66
commercial sports value, 103
homing tendency, 122
mark and recapture, 105–23
migration, 102–24
movements, seasonal, 102–24
tagging techniques, 104–5
Striped
blenny, 127
mullet, 255
Strobilation period, sea nettle, 211, 214
Stylochus ellipticus, 77, 78, 228

T

Tagatz, M. E., 202–3
Tagging, fish, 104–5
Talorchestia megalophthalma, 191
Tape grass, 15, 20
Taxonomy
methods, freshwater fishes, 30
lateral line system, cephalic canals, 30
Temperature
clam activity, 194–6
Cyathura, requirements, 14, 16, 20, 21, 25–6
epifauna, 74–5
interstitial waters, 19, 25–6
James River, 93, 94, 95, 99
MSX mortalities, 3
sea nettle, tolerance, 216–7
Tenellia, 81, 84
Terebra dislocata, 191
Thalassia, 15, 21
Thalassionema nitzschioides, 92, 100
Thalassiosira nordenskioldii, 92, 97, 100
rotula, 92, 93, 94, 100
subtilis, 92, 100
Thalassiothrix frauenfeldii, 100
Tiger shark, 53, 61
Toadfish, 86
Toxicity
of metal to fish, 262–3
salmonids, zinc, 263
zinc, 263
Trawl, 202
otter, 130
beam, 130
surface, 130–31
Trent, W. L., 130–1
Trinectes maculatus, 128
Tubeworms, 81–2, 86
Tube building, amphipod, 80, 81
Tunicates, 76–7, 78–9, 88
distribution, 79
Turbellarian, 78
Turbidity
epifauna, 75
estuary, measurements, 158
Turtle grass, 15

U

Uca, 22
Urothoe, 192

V

Vaillantia, 29, 30
description, 30
Vallisneria, 15, 16, 20, 21, 22, 23
Vernick, S. H. and V. Hearing, 170–86
Victorella pavida, 83
Villora, 30
Vision, shark, 57
Viviparus, 21–2
georgeianus, 23
Vorticellidae, 83

W

Warmouth bass, 22
Water
crowfoot, 224
zinc analyses, 262
Weakfish, 129
Weinstein, M. P. and R. Eisler, 253–8
White
perch, 66, 127, 135
antibody, 66
antibiotic sensitivity, 139
internal flora, 136, 143, 144–54
shark, 52, 53, 60, 61
tip shark, 53
White-winged doves, 201
Wilber, C. G. and R. Schneider, 63–5
Winter
flounder, 126–8
spawning patterns, 128

jellyfish, 219
salinity tolerance, 219
summer jellyfish similarities, 219
Wood, J. L. and J. D. Andrews, 1–13
Worm
flat, 78
gizzard, 68
neriid, 77
oligocharte, 21, 22, 23

Y

Yellow perch, 126, 128
spawning patterns, 128

Z

Zenaida asiatica, 201
Zenaidura macroura carolinensis, 200
Zinc, analyses, water, 262
toxicity to fish, 262, 263
Zoothamnium, 83

ERRATA

P. 52. Correct spelling: *"Carcharodon carcharias."*
P. 59. Correct spelling: *"Carcharhinus."*